

















Radial Velocity Studies of Close Binary Stars. III*

Slavek M. Rucinski, Wenxian Lu, Stefan W. Mochnacki

e-mail: *rucinski@astro.utoronto.ca*, *lu@astro.utoronto.ca*, *mochnacki@astro.utoronto.ca*

David Dunlap Observatory, University of Toronto

P.O.Box 360, Richmond Hill, Ontario L4C 4Y6, Canada

February 1, 2008

ABSTRACT

Radial velocity measurements and simple sine-curve fits to the orbital velocity variations are presented for the third set of ten contact binary systems: CN And, HV Aqr, AO Cam, YY CrB, FU Dra, RZ Dra, UX Eri, RT LMi, V753 Mon, OU Ser. All systems but two are contact, double-line spectroscopic binaries with four of them (YY CrB, FU Dra, V753 Mon, OU Ser) being the recent discoveries of the Hipparcos satellite project. The most interesting object is V753 Mon with the mass-ratio closest to unity among all contact systems ($q = 0.970 \pm 0.003$) and large total mass ($((M_1 + M_2)\sin^3 i = 2.93 \pm 0.06)$). Several of the studied systems are prime candidates for combined light and radial-velocity synthesis solutions.

Subject headings: stars: close binaries - stars: eclipsing binaries – stars: variable stars

1. INTRODUCTION

This paper is a continuation of the radial velocity studies of close binary stars, Lu & Rucinski 1999 (Paper I) and Rucinski & Lu 1999 (Paper II). The main goals and motivations are described in these papers. In short, we attempt to obtain modern radial velocity data for close binary systems which are accessible to the 1.8 meter class telescopes at medium spectral resolution of about $R = 10,000 - 15,000$. Selection of the objects is quasi-random in the sense that we started with shortest period contact binaries. The intention is to publish the results in groups of ten systems as soon as reasonable orbital elements can be obtained from measurements evenly distributed in orbital phases. We are currently observing a few dozens of such systems. The rate of progress may slow down, as we move into systems with progressively longer periods.

This paper is structured in the same way as Papers I and II in that it consists of two tables containing the radial velocity measurements (Table 1) and their sine-curve solutions (Table 2) and of brief summaries of previous studies for individual systems. The reader is referred to the

*Based on the data obtained at the David Dunlap Observatory, University of Toronto.

previous papers for technical details of the program. In short, all observations described here were made with the 1.88 meter telescope of the David Dunlap Observatory (DDO) of the University of Toronto. The Cassegrain spectrograph giving the scale of $0.2 \text{ \AA}/\text{pixel}$, or about $12 \text{ km/s}/\text{pixel}$, was used; the pixel size of the CCD was $19 \text{ }\mu\text{m}$. A relatively wide spectrograph slit of $300 \text{ }\mu\text{m}$ corresponded to the angular size on the sky of 1.8 arcsec and the projected width of 4 pixels. The spectra were centered at $5185 \text{ \AA}$ with the spectral coverage of $210 \text{ \AA}$. The exposure times were typically 10 – 15 minutes long.

The data in Table 1 are organized in the same manner as in Paper II. It provides information about the relation between the spectroscopically observed epoch of the primary-eclipse T_0 and the recent photometric determinations in the form of the (O–C) deviations for the number of elapsed periods E . It also contains, in the first column below the star name, our new spectral classifications of the program objects. The classification spectra, were obtained with a grating giving a dispersion of $0.62 \text{ \AA}/\text{pixel}$ in the range $3850 - 4450 \text{ \AA}$. The program-star spectra were “interpolated” between spectra of standard stars in terms of relative strengths of lines known as reliable classification criteria.

In the radial-velocity solutions of the orbits, the data have been assigned weights on the basis of our ability to resolve the components and to fit independent Gaussians to each of the broadening-function peaks. Weight equal to zero in Table 1 means that an observation was not used in our orbital solutions; however, these observations may be utilized in detailed modeling of broadening functions, if such are undertaken for the available material. The full-weight points are marked in the figures by filled symbols while half-weight points are marked by open symbols. Phases of the observations with zero weights are shown by short markers in the lower parts of the figures; they were usually obtained close to the phases of orbital conjunctions.

All systems discussed in this paper but one have been observed for radial velocities for the first time. The only exception is RZ Dra for which an SB1 orbit solution was obtained by Struve 1946. The solutions presented in Table 2 for the four circular-orbit parameters, γ , K_1 , K_2 and T_0 , have been obtained iteratively, with fixed values of the orbital period. First, two independent least-squares solutions for each star were made using the same programs as described in Papers I and II. Then, one combined solution for both amplitudes and the common γ was made with the fixed mean value of T_0 . Next, differential corrections for γ , K_1 , K_2 , and T_0 were determined, providing best values of the four parameters. These values are given in Table 2. The corrections to γ , K_1 , K_2 , and T_0 were finally subject to a “bootstrap” process (several thousand solutions with randomly drawn data with repetitions) to provide the median values and ranges of the parameters. We have adopted them as measures of uncertainty of parameters in Table 2.

Throughout the paper, when the errors are not written otherwise, we express standard mean errors in terms of the last quoted digits, e.g. the number $0.349(29)$ should be interpreted as 0.349 ± 0.029 .

2. RESULTS FOR INDIVIDUAL SYSTEMS

2.1. CN And

The variability of CN And was discovered by Hoffmeister 1949. Modern light curves were presented by Kaluzny 1983, Refert et al. 1985, Keskin 1989 and Samec et al. 1998. The light curve is of the EB-type, with unequally deep eclipses. From the eclipse timing, there exists a clear indication of a continuous period decrease. We used the almost contemporary with our observations determination of the primary eclipse time T_0 by Samec et al. 1998. The radial velocity curve of the secondary component (see Figure 1) shows some asymmetry in the first half of the orbital period which may explain why our T_0 is shifted by -0.014 day to earlier phases. From the radial-velocity point of view, the system looks like a typical A-type contact binary, but it can be also a very close semi-detached system.

Two groups of investigators attempted solutions of the light curves for orbital parameters, disregarding lack of any information on the spectroscopic mass-ratio. Our $q_{sp} = 0.39 \pm 0.03$ differs rather drastically from these photometric estimates. Kaluzny 1983 expected the mass-ratio to be within $0.55 < q_{ph} < 0.85$. Refert et al. 1985 found the most likely interval to be $0.5 < q_{ph} < 0.8$; they saw indications of a strong contact. The matter of the contact or semi-detached nature of the system should be re-visited in view of the discrepancy between our q_{sp} and the previous estimates of q_{ph} .

The photometric data at maximum light were published by Kaluzny 1983: $V = 9.62$, $(B - V) = 0.45$. The system is apparently a moderately strong X-ray source (Shaw et al. 1996).

2.2. HV Aqr

Variability of HV Aqr was discovered relatively recently by Hutton 1992. The type of variability was identified by Schirmer 1992 and a preliminary, but thorough study was presented by Robb 1992. The system shows total eclipses at the shallower of the two minima, so it is definitely an A-type one. This circumstance is important to our results because both available ephemerides of Schirmer and Robb do not predict the primary minimum correctly and we are not sure how our radial-velocity observations relate to the photometric observations: Our T_0 falls at phase 0.47 for the former and at 0.70 for the latter. Probably the values of the orbital period are slightly incorrect for both ephemerides. We used the period of 0.374460 day, following Schirmer 1992; if this is incorrect, then part of the scatter in our data may be due to the incorrect phasing over the span of 450 days of our observations.

The mass-ratio of HV Aqr is small, $q_{sp} = 0.145 \pm 0.05$. It agrees perfectly with the photometric solution of Robb 1992 of $q_{ph} = 0.146$ which confirms the validity of the photometric approach for totally eclipsing systems, in contrast with the typical lack of agreement for partially eclipsing systems.

Our spectral type of F5V agrees with the observed $(B - V) = 0.63 - 0.78$ for a relatively large reddening of $E_{B-V} = 0.08$ expected by Robb 1992. The system is bright with $V = 9.80$, and is potentially one of the best for a combined spectroscopic – photometric solution.

2.3. AO Cam

Variability of AO Cam was discovered by Hoffmeister 1966. Milone et al. 1982 analyzed their photometric observations in a simplified way. Subsequent photometric solution by Evans et al. 1985 and even the sophisticated work of Barone et al. 1993 did not bring much progress in view of the partial eclipses and total lack of any information on the mass-ratio. Barone et al. 1993 stated that AO Cam is definitely a W-type system; they estimated the photometric mass-ratio at $q_{ph} = 1.71 \pm 0.04$ or $1/q = 0.585$. This value is very different from our spectroscopic result, $q_{sp} = 0.413 \pm 0.011$, but we do confirm the W-type of the system.

AO Cam does not have any UBV data. Our spectral classification is G0V. In view of its considerable brightness of $V = 9.50$ at maxima, it appears to be a somewhat neglected system.

To predict the moment of the primary eclipse T_0 , we used the ephemeride based on the period of Evans et al. 1985 and the observations of the *secondary eclipses* by Faulkner 1986. The (O–C) deviation is relatively small in spite of many orbital periods elapsed since Faulker’s observations.

2.4. YY CrB

YY CrB is one of the variable stars discovered by the Hipparcos satellite mission (ESA 1997). The only available light curve comes from this satellite; the star has not been studied in any other way. There are no obvious indications of total eclipses, but the coverage of eclipses is relatively poor so that it is possible that the minimum identified as the primary is not the deeper one. We used the primary minimum ephemeride of Hipparcos and this results in a contact system of type A, that is with the more massive and hotter component eclipsed at this minimum.

The system is bright, $V_{max} = 8.64$, and its Hipparcos parallax is relatively large and well-determined, $p = 11.36 \pm 0.85$ milli-arcsec (mas), giving a good estimate of the absolute magnitude of the system, $M_V = 3.92 \pm 0.16$. With $(B - V) = 0.62 \pm 0.02$ from the Hipparcos database, the $M_V(\log P, B - V)$ calibration (Rucinski & Duerbeck 1997) gives $M_V^{cal} = 3.88$ so that the agreement is perfect. Our spectral classification of F8V agrees with the $(B - V)$ color index.

With the very good parallax and the new radial velocity data, YY CrB is one of the systems which have a great potential of providing an excellent combined photometric and spectroscopic solution.

2.5. FU Dra

FU Dra is another Hipparcos discovery. Again, we assumed the primary eclipse identification and its ephemeride as in the Hipparcos publication (ESA 1997). With these assumptions, the system appears to belong to the W-type systems. The mass-ratio is somewhat small for such systems, $q_{sp} = 0.25 \pm 0.03$.

The system is bright, $V_{max} = 10.55$, but its Hipparcos parallax is only moderately well determined, $p = 6.25 \pm 1.09$ mas, resulting in the absolute magnitude $M_V = 4.53 \pm 0.38$. The system was measured by the Hipparcos project to have a relatively large tangential motion, $\mu_{RA} = -255.85 \pm 1.18$ mas and $\mu_{dec} = 16.61 \pm 1.18$ mas. The large proper motion had been noticed before (Lee 1984a, Lee 1984b), but the spatial velocity was then estimated assuming an uncertain spectroscopic parallax. Using the Hipparcos data one obtains the two tangential components, $V_{RA} = -194$ km s⁻¹ and $V_{dec} = 13$ km s⁻¹. The radial velocity $\gamma = -11$ km s⁻¹ is moderate, so that only the RA component of the spatial tangential velocity is very large.

2.6. RZ Dra

RZ Dra has been frequently photometrically observed since the discovery by Ceraski 1907. The most recent extensive analysis of the system was by Kreiner et al. 1994. It utilized the only extant set of spectroscopic observations by Struve 1946 which led to a detection of one, brighter component. Our data confirm the primary amplitude $K_1 = 100$ km s⁻¹, but we have been to also detect the secondary component. The system consists of components considerably differing in the effective temperature, and thus is classified as an EB-type binary. Spectroscopically, one sees a more massive component eclipsed at the deeper minimum so that it can be an Algol semi-detached binary or an A-type contact system in poor thermal contact. The analysis of Kreiner et al. 1994 was made under the assumption of the semi-detached configuration. They found the photometric mass-ratio $q_{ph} \simeq 0.45$, but some solutions suggested $q_{ph} \simeq 0.55$. Our relatively well defined solution for both components gives $q_{sp} = 0.40 \pm 0.04$. The same investigation of Kreiner et al. provided the starting value of T_0 . In spite of indications that the period may be variable and that many epochs elapsed since the study by Kreiner et al., the observed shift in the primary eclipse time is relatively small.

The spectral type that we observed, A6V, most probably applies to the primary component which is much hotter than its companion (we have not attempted to separate the spectra in terms of spectral types). The Hipparcos parallax of the system, $p = 1.81 \pm 1.01$ mas, is too poor for an more extensive analysis of the absolute magnitude of the system. RZ Dra appears to be a relatively short-period (0.55 day) semi-detached Algol with both components accessible to spectroscopic observations.

2.7. UX Eri

UX Eri is a contact binary which was extensively photometrically observed since its discovery by Soloviev 1937. The first modern contact-model solutions which gave surprisingly good agreement with our spectroscopic mass-ratio, $q_{sp} = 0.37 \pm 0.02$, was presented by Mauder 1972 almost quarter of a century ago; it utilized the light curve of Binnendijk 1967 and arrived at $q_{ph} = 0.42$.

Our observations have been supplemented by four observations obtained at the same time by Dr. Hilmar Duerbeck at the European Southern Observatory with a 1.52m telescope and a Cassegrain spectrograph. As a starting point of our solution, we used the moment of the primary minimum T_0 as predicted on the basis of observations by Agerer & Huebscher 1998a and Agerer & Huebscher 1998b; both took place actually slightly after our spectroscopic observations.

UX Eri appears to be an A-type contact binary. The $(B - V)$ color index is not available for this star (it has also not been measured by Hipparcos), so we have not been able to relate it to our spectral classification of F9V. The Hipparcos parallax $p = 6.57 \pm 2.84$ mas provides a relatively poor estimate of the absolute magnitude for the maximum brightness of $V_{max} = 10.59$: $M_V = 4.7 \pm 0.9$.

2.8. RT LMi

RT LMi was discovered as a variable star by Hoffmeister 1949. The most recent analysis from which we took the time of the primary eclipse T_0 is by Niarchos et al. 1994. The system has been characterized in this study as a W-type contact binary of spectral type G0V with photospheric spots. Our spectral classification is based on poor spectra, but they indicate a slightly earlier spectral type, of approximately F7V. As Niarchos et al. 1994 pointed out, the minima were observed to be of almost equal depth. Our spectroscopic orbit gives an A-type so that the minimum selected by the authors as the primary corresponds to the eclipse of the more massive component (the temporal shift is very small in spite of many elapsed epochs). The photometric solution of Niarchos et al. 1994 assuming the W-type appears therefore to be invalid. The system is otherwise quite an inconspicuous contact binary. It lacks even most essential photometric data. The Simbad database gives $V_{max} = 11.4$, but the source of this value is not cited.

2.9. V753 Mon

V753 Mon is a new discovery of the Hipparcos mission. It is probably one of the most interesting new close binaries recently discovered. V753 Mon has not been studied before. The only published photometric data come from the *uvby* survey of Olsen 1994 who found $(b - y) = 0.214 \pm 0.008$, $m_1 = 0.160 \pm 0.003$ and $c_1 = 0.693 \pm 0.023$. The $(b - y)$ color index color

corresponds to $(B - V) \simeq 0.34$ or the spectral type F2V. Our spectral classification is A8V which does not agree with these estimates and with $(B - V) = 0.36$ in the Hipparcos catalog, unless there is considerable reddening of about $E_{B-V} \simeq 0.12$. However, the early type would be in a better accord with the large masses indicated by the radial velocity solution (see below). The brightness data in the Olsen measurements indicated quite appreciable variability, $V = 8.46 \pm 0.34$, but this indication has been apparently overlooked. The Hipparcos mission database treats it as a new discovery.

The two features distinguish V753 Mon as a particularly interesting system: The mass-ratio close to unity and the large amplitudes of radial velocity variations indicating a large total mass. The mass-ratio $q_{sp} = 0.970 \pm 0.009$ is the closest to unity of all known contact binaries. Note that the currently largest mass ratios are $q = 0.80$ for VZ Psc (Hrivnak et al. 1995) and SW Lac (Zhai & Lu 1989) and $q = 0.84$ for OO Aql (Hrivnak 1989). Since contact systems with $q \simeq 1$ are not observed, but are expected to experience *strong favorable observational biases for their detection and ease in analysis*, it is generally thought that contact configurations avoid this particular mass-ratio. The summed amplitudes of the radial velocity variations for V753 Mon give the total mass of the system, $(M_1 + M_2)\sin^3 i = 2.93 \pm 0.06$. This is in perfect agreement with the expected masses of two main-sequence stars of the spectral type F2V seen on an orbit exactly perpendicular to the plane of the sky. The light variation is about 0.52 mag., in place of the expected about 1.0 mag. for a contact system with $q \simeq 1$; therefore, the total mass for $i < 90^\circ$ may turn out to be substantially larger. For the spectral type estimated by us the individual masses should be close to $1.7 M_\odot$.

Our radial velocity data show that with the ephemeride based on the Hipparcos data, the system belongs to the W-type contact systems. Apparently, the eclipses are of almost the same depth, as expected for $q \simeq 1$. However, the light curve from Hipparcos is poorly covered around the secondary minimum so that the identification of the eclipses is uncertain. Obviously, the distinction between A-type and W-type systems becomes immaterial for $q \rightarrow 1$.

The system is begging a new light curve and an extensive analysis, not only because of the unusual properties, but also because it is bright, $V_{max} = 8.34$, and has a moderately well determined Hipparcos parallax: $p = 5.23 \pm 1.04$ resulting in $M_V = 1.93 \pm 0.43$. This is in perfect agreement with the $M_V(\log P, B - V)$ calibration which gives $M_V^{cal} = 1.90$ for the assumed $(B - V) = 0.34$. The agreement would not be that good if the color index is smaller, say 0.22, then one would obtain $M_V^{cal} = 1.54$ which is still within the uncertainty of the parallax. Further investigations of V753 Mon will therefore contribute to the absolute-magnitude calibration which is rather moderately-well defined for contact binaries with periods longer than about 0.5 day (Rucinski & Duerbeck 1997).

2.10. OU Ser

OU Ser is the fourth Hipparcos mission discovery in this group of systems. The light curve shows almost equally deep minima. With the Hipparcos ephemeride, the system appears to be an A-type one with a small mass-ratio $q_{sp} = 0.173 \pm 0.017$. Our spectral classification of the system indicates the spectral type F9/G0V.

The distinguishing properties of OU Ser in the Hipparcos database are its large proper motion and a well measured parallax. The tangential components of the proper motion are $\mu_{RA} = -387.5 \pm 0.9$ mas and $\mu_{dec} = 2.8 \pm 0.8$ mas. With the parallax of $p = 17.3 \pm 1.0$ mas this translates into the spatial components $V_{RA} = -106$ km s⁻¹ and $V_{dec} = 1$ km s⁻¹. The mean radial velocity of the system is $\gamma = -64.08 \pm 0.41$ km s⁻¹. Thus, the RA and the radial velocity components indicate a high-velocity star.

The large proper motion of the star had been the reason for inclusion in the survey by Carney et al. 1994. They noted also the broad lines indicating short-period binarity and possibility of light variations. Their photometric data $V = 8.27$, $(B - V) = 0.62$ and $(U - B) = 0.08$ agree with our spectral classification, F9/G0V. The *ubvy* survey of Olsen 1994 suggests a slightly larger $(B - V) \simeq 0.66$ on the basis of the $(b - y) = 0.411 \pm 0.003$, hence a spectral type around G1/2V. The difference in the classification may be due to the apparently low metallicity of the system as judged by its low index $m_1 = 0.168 \pm 0.004$ (provided this index is not confused by any chromospheric activity). the other data of Olsen are $V = 8.278 \pm 0.005$ and $c_1 = 0.281 \pm 0.006$.

Assuming $V_{max} = 8.25$ and the parallax $p = 17.3 \pm 1.0$ mas, one obtains $M_V = 4.44 \pm 0.12$. This again agrees very well with the absolute magnitude derived from the $M_V(\log P, B - V)$ calibration of Rucinski & Duerbeck 1997: For $(B - V) = 0.62$, $M_V^{cal} = 4.32$, while for $(B - V) = 0.66$, $M_V^{cal} = 4.46$.

With the excellent parallax data and its properties of a high velocity star, OU Ser deserves a combined photometric – spectroscopic solution.

3. SUMMARY

The paper brings radial velocity data for the third group of ten close binary systems that we observed at the David Dunlap Observatory. All, but RZ Dra (which had SB1 radial velocity data), have never been observed spectroscopically; all ten are binaries with both components clearly detected so that they can be called SB2. All systems, but CN And and RZ Dra, which may be very close semi-detached systems, are contact binaries. We describe special features of the individual systems in the descriptions in Section 2. We note that again about half of the system are A-type contact binaries; the likely reasons why we prefer them over the W-type systems in our randomly drawn sample are given in the Conclusions to Paper II. We also observed as EB2 systems two binaries, CN And and RZ Dra; they are most probably semi-detached binaries.

We do not give the calculated values of $(M_1 + M_2) \sin^3 i = 1.0385 \times 10^{-7} (K_1 + K_2)^3 P(\text{day}) M_\odot$ because in most cases the inclination angles are either unknown or not trustworthy. However, one case is very interesting here: The total mass of components of the system V753 Mon is very large, $M_1 + M_2 > 2.93 M_\odot$. Since *too large velocity amplitudes* are a rare phenomenon in the world of contact systems, this system requires special attention of the observers. The binary is also unique in having its mass-ratio exceptionally close to unity, $q_{sp} = 0.970 \pm 0.009$. Two other systems discovered by the Hipparcos mission are also very important and promise excellent combined solutions, YY CrB and OU Ser. It is important that both are high-velocity stars and have excellent parallaxes. And, finally, the recently discovered system HV Aqr offers an excellent solution in view of the total, well-defined eclipses and very good radial velocity data.

The authors would like to thank Jim Thomson for help with observations and to Hilmar Duerbeck for a permission to use his observations of UX Eri.

The research has made use of the SIMBAD database, operated at the CDS, Strasbourg, France and accessible through the Canadian Astronomy Data Centre, which is operated by the Herzberg Institute of Astrophysics, National Research Council of Canada.

REFERENCES

- Inf. Bull. Var. Stars, 4562
- Inf. Bull. Var. Stars, 4606
- Barone, F., Di Fioere, L., Milano, L. & Russo, G. 1993, ApJ, 407, 237
- Binnendijk, L. 1967, AJ, 72, 82
- Carney, B.W., Latham, D.W., Laird, J.B. & Aguilar, L.A. 1994, AJ, 107, 2240
- Ceraski, W. 1907, Astr. Nachr., 174, 265
- European Space Agency. 1997. The Hipparcos and Tycho Catalogues (Paris: ESA), SP-1200
- Evans III, E., Grosseohme, D.H. & Moyer Jr., E.J. PASP, 97, 648
- Falkner, D.R. 1986, PASP, 98, 690
- Hoffmeister, C. 1949, Astr. Nachr., 12, 1
- Hoffmeister, C. 1949b, Astr. Abhandl., 12, 1
- Hoffmeister, C. 1966, Astr. Nachr., 289, 1
- Hrivnak, B.J. 1989, ApJ, 340, 458

- Hrivnak, B.J., Guinan, E.F. & Lu, W. 1995, *ApJ*, 455, 300
- Hutton, R.G. 1992, *Inf. Bull. Var. Stars*, 3723
- Kaluzny, J. 1983, *Acta Astr.*, 33, 345
- Keskin, V. 1989, *Ap&SS*, 153, 191
- Kreiner, J.M., Pajdosz, G., Tremko, J. & Zola, S. 1994, *A&A*, 285, 459
- Lee, S.-G. 1984a, *AJ*, 89, 702
- Lee, S.-G. 1984b, *AJ*, 89, 720
- Lu, W. & Rucinski, S.M. 1999, *AJ*, 118, 515 (Paper I)
- Mauder, H. 1972, *A&A*, 17, 1
- Milone, E.F., Piggott, D.H. & Morris, S.L. *JRASC*, 76, 90
- Niarchos, P.G., Hoffmann, M. & Duerbeck, H.W. 1994, *A&AS*, 103, 39
- Olsen, E.H. 1994, *A&AS*, 106, 257
- Rafert, J.B., Markworth, N.L. & Michaels, E.J. 1985 *PASP*, 97, 310
- Robb, R.M. 1992, *Inf. Bull. Var. Stars*, 3798
- Rucinski, S.M. & Duerbeck, H.W. 1997, *PASP*, 109, 1340
- Rucinski, S.M. & Lu, W. 1999, *AJ*, 118, 2451 (Paper II)
- Samec, R.G., Laird, H., Mutzke, M. & Faulkner, D.R. *Inf. Bull. Var. Stars*, 4616
- Shaw, J.S., Caillault, J., Schmitt, J.H.M.M. 1996, *ApJ*, 461, 951
- Shirmer, J. 1992, *Inf. Bull. Var. Stars*, 3785
- Soloviev, A. 1937, *Tadjik Obs. Circ.*, No. 27, 7
- Struve, O. 1946, *ApJ*, 103, 76
- Zhai, D.S. & Lu, W.X. 1989, *Acta Astr. Sinica*, 9, 208

Captions to figures:

Fig. 1.— Radial velocities of the systems CN And, HV Aqr, AO Cam and YY CrB are plotted in individual panels versus orbital phases. The thin lines give the respective circular-orbit (sine-curve) fits to the radial velocities. Note that AO Cam is a W-type system while the rest are A-type systems. Short marks in the lower parts of the panels show phases of available observations which were not used in the solutions because of the blending of lines. Open symbols in this and the next figure indicate observations given half-weights in the solutions. All panels have the same vertical scales.

Fig. 2.— Radial velocities of the systems FU Dra, RZ Dra, UX Eri and RT LMi.

Fig. 3.— Radial velocities of the systems V753 Mon and OU Ser. While both of these recently-discovered systems are very interesting, V753 Mon is exceptional in having a mass-ratio very close to unity in showing very large radial-velocity amplitudes.

TABLE 1. DDO observations of the third group of ten close binary systems

HJD-2,400,000	Phase	V_1	ΔV_1	V_2	ΔV_2
CN And					
50615.7873	0.3120	-107.6	-1.8	178.0	-4.7
50615.7986	0.3364	-105.5	-5.7	154.6	-12.6
50615.8112	0.3636	-91.3	-0.3	133.3	-11.4
50615.8222	0.3874	-73.2	8.6	92.0	-28.9
50615.8339	0.4127	-52.5	18.0	...	...
50631.8015	0.9154	40.6 ^a	21.1 ^a	...	...
50631.8122	0.9385	...	...	...	...
50631.8250	0.9662	...	...	...	...
50651.7711	0.0656	...	...	...	...
50651.7821	0.0893	-76.6 ^a	-5.1 ^a	...	...
50651.7948	0.1168	-80.4	3.1	156.6	31.2
50651.8056	0.1401	-91.2	1.1	169.0	20.9
50651.8173	0.1654	-98.1	2.2	170.1	1.6
50651.8282	0.1889	-106.1	-0.1	184.7	1.5
50651.8416	0.2179	-109.2	1.4	195.0	0.1
50651.8525	0.2414	-106.8	5.5	198.0	-1.2
50651.8648	0.2680	-103.4	8.4	185.5	-12.5
50700.7094	0.8111	55.0	-1.3	-250.5	-17.6
50700.7206	0.8353	57.4	7.0	-250.5	-32.7
50700.7326	0.8612	51.4	9.3	-225.6	-29.0
50700.7433	0.8844	42.3	9.0	-205.4	-31.5
50708.7686	0.2254	-103.2	8.2	203.6	6.8
50708.7829	0.2563	-102.4	9.9	206.7	7.4
50708.7952	0.2828	-118.6	-8.1	191.7	-3.0
50709.7059	0.2507	-113.3	-0.9	186.1	-13.4
50709.7181	0.2770	-109.7	1.4	186.6	-9.6
50771.5187	0.8154	59.7	4.3	-244.8	-14.2
50771.5326	0.8454	49.9	2.5	-213.8	-3.7
50852.4794	0.7547	56.3	-6.3	-262.5	-13.4
50852.4904	0.7785	61.9	0.6	-232.7	12.9
50852.5032	0.8061	58.9	1.6	-247.0	-11.6
51080.6489	0.7818	60.4	-0.5	-216.1	28.6
51080.6597	0.8052	62.2	4.7	-206.1	29.8
51080.6718	0.8313	52.3	0.8	-220.5	0.1
51080.6826	0.8547	46.0	1.6	-206.0	-3.6
51080.6943	0.8799	34.3	-0.8	-205.6	-27.1
51080.7054	0.9039	31.5	6.7	-180.5	-28.3
51085.6730	0.6379	36.2	-5.6	-208.0	-12.2
51085.6842	0.6621	58.2	8.6	-227.6	-11.8
51085.6959	0.6874	58.3	2.3	-225.1	7.0
51085.7065	0.7103	61.5	1.6	-239.8	2.5
51085.7208	0.7412	67.1	4.6	-228.5	20.4
51085.7323	0.7660	59.0	-3.2	-237.9	10.2
51085.7460	0.7956	61.8	2.7	-222.1	18.0
51085.7573	0.8200	49.8	-4.5	-193.6	34.2

TABLE 1. (continued)

HJD-2,400,000	Phase	V ₁	ΔV_1	V ₂	ΔV_2
HV Aqr					
50631.7429	0.1851	-33.4	9.3	264.9	-0.7
50785.5092	0.8199	31.9	-2.7	-273.0	-3.7
50785.5204	0.8498	31.0	0.4	-252.4	-11.1
51080.5304	0.6776	32.7	-1.6	-270.5	-3.2
51080.5416	0.7075	32.5	-4.6	-278.6	8.1
51080.5542	0.7411	37.0	-1.6	-294.0	2.7
51080.5650	0.7700	37.6	-0.7	-297.5	-2.7
51080.5768	0.8015	30.9	-5.5	-271.9	10.0
51080.5875	0.8301	34.3	0.9	-263.6	-2.8
51080.5992	0.8613	29.6	0.9	-224.9	3.3
51080.6099	0.8899	30.8	7.5	-187.8	3.1
51080.6216	0.9211	...	...	...	...
51080.6323	0.9497	...	...	...	...
51085.5773	0.1554	-36.7	2.2	237.8	-1.5
51085.5880	0.1840	-44.5	-1.9	279.6	14.9
51085.5998	0.2155	-50.8	-5.6	287.2	4.5
51085.6104	0.2438	-53.3	-7.2	288.3	-1.1
51085.6235	0.2788	-44.7	0.8	267.2	-17.6
51085.6342	0.3073	-35.9	7.6	281.9	11.2
51085.6459	0.3386	-49.1	-9.3	257.7	12.4
51085.6566	0.3672	-42.7	-7.5	206.0	-7.6
AO Cam					
50699.8482	0.0913	-160.6 ^a	-11.5 ^a	42.4	-0.6
50699.8589	0.1237	-190.4	-1.5	70.0	10.6
50699.8725	0.1649	-233.5	-4.7	84.2	8.3
50699.8832	0.1974	-250.7	-0.6	77.4	-7.3
50699.8952	0.2338	-263.0	-0.6	93.5	3.7
50850.4835	0.6933	216.3	-5.3	-110.6	-0.4
50850.4964	0.7324	236.8	1.0	-111.6	4.5
50850.5111	0.7770	234.7	1.0	-110.6	4.6
50850.5244	0.8173	216.0	0.7	-106.9	0.7
50850.5387	0.8607	182.7	3.5	-89.1	3.6
50850.6490	0.1950	-257.0	-8.1	82.8	-1.4
50850.6629	0.2371	-260.9	2.0	89.6	-0.4
50850.6770	0.2799	-255.4	3.9	80.4	-8.1
50850.6889	0.3159	-244.1	-1.6	79.6	-2.0
50850.7030	0.3587	-201.3	6.2	74.0	6.9
50850.7158	0.3975	-170.3	-6.7	50.7	1.7
50850.7303	0.4414	...	...	...	...
50852.5208	0.8687	182.3	11.5	-88.2	1.0
50852.5294	0.8948	159.0 ^a	18.4 ^a	-72.1	4.6
50852.5394	0.9251	...	...	-55.0 ^a	5.1 ^a
51220.6274	0.6645	203.6	1.6	-97.0	5.1
51220.6382	0.6972	218.5	-5.2	-123.1	-12.1
51220.6499	0.7327	233.8	-2.1	-120.5	-4.4
51220.6606	0.7651	228.6	-7.6	-107.3	8.9
51220.6735	0.8042	218.9	-4.0	-114.2	-3.5
51220.6845	0.8376	195.9	-4.5	-106.9	-5.5

TABLE 1. (continued)

HJD-2,400,000	Phase	V ₁	ΔV_1	V ₂	ΔV_2
YY CrB					
50948.6816	0.9077	50.8	18.1	-149.6	8.3
50948.6888	0.9268	42.5	16.9	-123.6	5.1
50948.6960	0.9460	42.6	24.5	-114.7	-16.9
50948.7122	0.9890	...	...	...	...
50948.7202	0.0102	...	...	...	...
50948.7277	0.0302	...	...	...	...
50949.6499	0.4791	...	...	...	...
50949.6571	0.4983	...	...	...	...
50949.6643	0.5174	...	...	...	...
50951.6721	0.8493	57.3	6.6	-232.6	-0.8
50951.6797	0.8694	60.4	15.2	-187.5	21.8
50951.6872	0.8894	42.4	3.4	-195.0	-11.2
50951.6959	0.9125	46.7	15.7	-152.4	-1.5
50951.7066	0.9409	42.2	22.1	-139.6	-33.4
50960.7595	0.9816	...	...	...	...
50960.7676	0.0031	...	...	...	...
50960.7751	0.0230	...	...	...	...
50960.7834	0.0451	...	...	...	...
50960.7907	0.0645	...	...	...	...
50960.7979	0.0836	...	...	...	...
50960.8066	0.1067	-55.3	-8.4	183.1	13.8
50960.8138	0.1258	-60.1	-7.1	209.1	14.8
50960.8210	0.1449	-64.8	-6.4	217.5	1.0
50960.8290	0.1662	-65.5	-2.1	234.3	-3.0
50960.8362	0.1853	-67.6	-0.5	255.0	2.5
50960.8434	0.2044	-71.9	-2.0	257.4	-6.5
50960.8514	0.2257	-68.2	3.7	270.6	-1.4
50961.6013	0.2171	-71.2	0.0	269.9	0.6
50961.6088	0.2370	-67.8	4.6	274.2	-0.1
50961.6160	0.2561	-75.8	-3.2	265.5	-9.6
50961.6249	0.2797	-67.9	3.6	262.9	-7.5
50961.6321	0.2989	-63.9	5.6	264.0	1.8
50961.6393	0.3180	-64.5	2.0	240.2	-9.9
50961.6476	0.3400	-65.3	-3.2	233.2	1.5
50961.6549	0.3594	-55.6	1.6	218.9	7.2
50961.6620	0.3783	-54.2	-2.5	191.7	2.5
50961.6704	0.4006	-48.3	-3.9	166.4	7.3
50961.6776	0.4197	-41.5	-4.0	135.8	5.1
50961.7762	0.6815	49.2	-8.1	-271.4	-12.5
50961.7834	0.7007	53.3	-6.9	-277.7	-6.6
50961.7906	0.7198	56.4	-5.9	-280.0	-0.6
50961.7994	0.7431	49.7	-13.7	-294.2	-10.0
50961.8066	0.7623	53.5	-9.8	-287.8	-4.2
50961.8138	0.7814	53.2	-9.0	-278.5	0.5
50961.8228	0.8053	51.6	-7.8	-269.4	-1.7
50961.8300	0.8244	54.4	-1.8	-254.2	0.2
50961.8372	0.8435	57.4	5.3	-235.1	2.4

TABLE 1. (continued)

HJD-2,400,000	Phase	V ₁	ΔV_1	V ₂	ΔV_2
FU Dra					
50853.7375	0.1173	-208.4	-8.3	40.0	4.1
50853.7475	0.1499	-228.6	9.8	56.4	10.9
50853.7583	0.1851	-269.6	-0.5	48.7	-4.5
50853.7679	0.2164	-294.7	-8.8	47.4	-10.0
50853.7792	0.2532	-284.4	7.7	55.1	-3.9
50853.7889	0.2849	-288.6	-3.2	52.8	-4.5
50853.8002	0.3217	-263.3	0.8	58.4	6.4
50853.8868	0.6040	170.7	11.3	-53.1	1.1
50853.8978	0.6399	201.4	-3.4	-73.7	-8.1
50853.9104	0.6810	240.3	-3.1	-77.9	-2.6
50853.9204	0.7136	266.9	4.8	-80.9	-1.0
50853.9311	0.7485	268.3	-1.1	-80.2	1.6
50853.9409	0.7804	271.0	6.7	-72.9	7.6
50853.9521	0.8169	257.7	12.8	-72.9	2.7
50858.8237	0.6999	255.2	-0.4	-71.6	6.7
50858.8238	0.7003	241.0	-14.8	-90.7	-12.3
50858.8482	0.7798	268.2	3.7	-79.2	1.3
50858.8595	0.8167	227.4	-17.7	-90.9	-15.2
50860.8036	0.1551	-240.6	3.0	52.8	6.0
50860.8313	0.2454	-277.0	15.0	54.9	-4.1
50860.8582	0.3331	-263.0	-8.2	42.7	-6.9
50860.8882	0.4309	...	...	...	...
50896.8648	0.7262	274.1	7.9	-66.3	14.7
50880.9355	0.7916	265.5	5.7	-89.0	-9.6
RZ Dra					
50610.8110	0.2287	-74.3	11.0	239.2	-13.8
50610.8229	0.2503	-90.6	-4.4	258.0	2.8
50610.8360	0.2741	-74.6	10.5	251.2	-1.2
50645.6647	0.4983	...	...	...	...
50645.6794	0.5250	...	...	...	...
50645.6954	0.5540	...	...	...	...
50700.5321	0.0986	-53.3 ^a	-7.8 ^a	...	...
50700.5464	0.1245	-51.2	6.4	184.8	1.8
50700.5620	0.1528	-61.4	7.3	209.1	-1.9
50700.5764	0.1790	-80.1	-3.4	212.5	-18.7
50700.5917	0.2068	-70.0	12.6	243.3	-2.9
50700.6059	0.2325	-78.1	7.5	248.4	-5.3
50700.6217	0.2612	-84.8	1.1	255.8	1.3
50700.6368	0.2886	-92.1	-8.8	226.6	-21.4
50700.6539	0.3197	-74.7	2.3	229.5	-2.6
50700.6686	0.3464	-73.6	-4.6	231.4	19.7
50700.6854	0.3768	-53.4	3.6	216.4	34.9
50708.5343	0.6249	71.8	-7.3	-187.3	-25.2
50708.5485	0.6507	95.9	6.6	-213.7	-26.0
50708.5644	0.6795	89.2	-9.0	-169.8	40.4
50708.5787	0.7055	96.9	-6.9	-232.7	-8.4
50708.5940	0.7333	114.0	7.0	-229.8	2.6
50708.6083	0.7592	108.9	1.5	-189.6	43.8
50708.6242	0.7881	100.6	-4.2	-214.4	12.4
50708.6389	0.8148	118.3	18.7	-223.0	-9.2
50708.6519	0.8384	...	...	...	...

TABLE 1. (continued)

HJD-2,400,000	Phase	V ₁	ΔV_1	V ₂	ΔV_2
51326.6143	0.6196	69.6	-7.2	-180.9	-24.6
51326.6267	0.6421	73.9	-12.2	-226.0	-46.2
51326.6401	0.6665	94.0	-0.5	-218.2	-17.3
51326.6543	0.6922	99.7	-1.5	-216.8	1.1
51326.6675	0.7162	108.1	2.7	-233.0	-4.7
51326.6798	0.7385	112.9	5.6	-230.2	3.0
51326.6918	0.7603	108.2	0.9	-244.2	-10.9
51326.7049	0.7841	120.5	15.2	-231.5	-3.3
51326.7174	0.8068	102.7	1.3	-201.8	16.6
51326.7318	0.8329	109.8	15.1	-221.1	-19.7
51326.7437	0.8545	99.9	12.5	-186.4	-3.5
51459.5493	0.9353	71.3 ^a	22.3 ^a	...	...
UX Eri					
50414.6722	0.7633	110.2	6.0	-241.0	-8.9
50414.6808	0.7826	112.6	10.0	-220.5	7.3
50415.5345 ^b	0.6998	107.6	7.6	-208.1	12.8
50416.6548 ^b	0.2158	-78.0	-1.1	245.0	-7.9
50417.6084 ^b	0.3574	-55.6	3.3	209.4	4.7
50418.7036 ^b	0.8169	94.9	-1.6	-204.8	6.8
50470.4851	0.1069	-46.6	-2.3	188.7	22.9
50470.4957	0.1308	-58.2	-3.8	206.2	13.5
50470.5084	0.1593	-66.6	-2.1	217.5	-2.2
50470.5198	0.1849	-76.2	-4.8	242.1	3.8
50470.5325	0.2134	-83.7	-7.1	250.9	-1.2
50470.5432	0.2374	-79.1	-0.4	259.8	2.0
50700.7652	0.2661	-93.6	-15.1	247.7	-9.6
50700.7794	0.2980	-70.2	4.6	246.6	-0.9
50700.7947	0.3323	-69.8	-2.8	226.7	0.3
50700.8089	0.3642	-52.2	4.1	208.4	10.5
50700.8251	0.4006	-43.0	-2.1	180.4 ^a	23.9 ^a
50709.8225	0.6068	61.8	-8.0	-159.0 ^a	-19.0 ^a
50709.8353	0.6356	89.9	8.1	-175.7	-3.6
50854.5324	0.5938	54.5	-9.3	-146.2 ^a	-22.3 ^a
50854.5454	0.6230	80.9	4.0	-161.9	-3.1
50854.5584	0.6522	73.5	-14.3	-175.6	12.4
51229.4903	0.6679	83.1	-9.5	-187.5	13.5
51229.5010	0.6919	92.6	-5.9	-205.9	10.9
51229.5128	0.7184	96.9	-5.8	-226.9	1.3
51229.5237	0.7429	100.5	-4.0	-231.6	1.1
51229.5361	0.7708	95.8	-8.0	-240.6	-9.7
51229.5468	0.7948	96.3	-4.6	-230.6	-7.3
RT LMi					
51254.5913	0.6163	44.1	-9.3	-206.1	-22.1
51254.6031	0.6478	64.7	-1.4	-234.4	-15.7
51254.6169	0.6846	66.4	-10.7	-257.2	-8.4
51254.6287	0.7161	91.2	8.3	-238.4	26.2
51254.6408	0.7484	96.9	11.9	-241.3	29.2
51254.6549	0.7860	68.7	-13.9	-266.7	-2.8
51254.6668	0.8177	81.4	4.9	-247.6	-0.3
51254.6796	0.8519	76.9	10.7	-222.2	-3.2
51254.6916	0.8839	58.9	5.6	-187.3	-3.6
51261.5672	0.2228	-102.7	1.5	246.0	-0.1

TABLE 1. (continued)

HJD-2,400,000	Phase	V ₁	ΔV_1	V ₂	ΔV_2
51261.5790	0.2543	-111.6	-6.0	255.9	6.1
51261.5919	0.2887	-95.5	7.3	249.7	7.5
51261.6038	0.3204	-91.5	5.0	213.5	-11.3
51261.7384	0.6794	70.8	-5.0	-264.1	-18.8
51261.7507	0.7122	90.7	8.3	-232.4	30.8
51261.7638	0.7472	80.2	-4.8	-281.1	-10.7
51261.7761	0.7800	77.6	-5.8	-275.1	-9.2
51267.5473	0.1732	-100.9	-6.1	206.2	-14.0
51267.5584	0.2028	-100.1	1.4	262.7	24.2
51267.5749	0.2468	-112.0	-6.4	238.2	-11.6
51267.5892	0.2850	-100.0	3.3	232.6	-11.0
51267.6009	0.3162	-101.2	-3.7	220.8	-6.9
51267.6145	0.3525	-81.2	5.4	199.8	2.0
51267.6252	0.3810	-74.6	0.5	193.4	26.8
51267.6382	0.4157	-74.1 ^a	-15.6 ^a	...	...
51267.6494	0.4456	...	...	...	...
51267.6608	0.4760	...	...	...	...
V753 Mon					
50852.6964	0.6378	163.6	-9.1	-103.6	-12.1
50852.7109	0.6593	178.6	-8.2	-112.7	-7.5
50852.7409	0.7036	207.5	0.3	-124.7	0.3
50852.7497	0.7166	204.1	-6.7	-126.9	1.5
50852.7593	0.7308	210.3	-3.1	-126.0	4.9
50852.7681	0.7437	216.9	2.4	-123.7	8.3
50854.5887	0.4328	...	...	...	...
50854.5984	0.4471	...	...	...	...
50858.5422	0.2721	-127.3	8.5	219.6	11.9
50858.5518	0.2863	-124.9	8.0	206.3	1.4
50858.5631	0.3030	-123.0	4.8	205.8	5.8
50858.5715	0.3154	-117.6	5.2	207.1	12.0
50884.5606	0.7012	200.9	-5.5	-123.0	1.2
50884.5718	0.7177	202.4	-8.7	-126.7	2.0
50884.5857	0.7383	214.9	0.7	-122.7	9.0
50884.5980	0.7564	214.7	0.2	-126.3	5.7
51232.5375	0.6624	192.6	4.0	-112.6	-5.7
51232.5481	0.6781	193.8	-3.2	-118.9	-3.9
51232.5600	0.6956	204.6	0.1	-124.3	-2.0
51232.5716	0.7128	206.8	-3.1	-126.1	1.4
51232.5856	0.7334	213.8	0.1	-131.6	-0.4
51232.5971	0.7504	212.5	-2.2	-129.7	2.5
51232.6111	0.7711	213.7	0.6	-130.5	0.2
51232.6226	0.7881	208.1	-1.5	-125.1	2.2
51232.6370	0.8094	207.1	4.5	-119.3	1.1
51232.6476	0.8250	200.0	4.5	-112.3	1.3
51232.6601	0.8435	193.4	8.3	-99.0	4.6
51232.6707	0.8591	181.2	6.3	-93.4	0.2
51232.6840	0.8788	168.4	8.3	-77.5	1.8
51232.6949	0.8949	158.4	11.8	-64.0	2.2
51281.5742	0.0895	-80.9 ^a	-25.6 ^a	123.3	-6.3
51281.5856	0.1063	-92.4	-21.9	134.6	-9.8
51281.5972	0.1235	-99.8	-15.1	151.1	-7.1
51513.8070	0.0969	-92.5 ^a	-30.4 ^a	126.7	-9.5

TABLE 1. (continued)

HJD-2,400,000	Phase	V ₁	ΔV_1	V ₂	ΔV_2
51513.8180	0.1131	-103.2	-26.9	140.3	-9.7
51513.8304	0.1315	-109.1	-18.2	159.6	-4.5
51513.8412	0.1474	-110.6	-8.5	174.3	-0.8
51513.8536	0.1657	-123.4	-10.0	184.1	-1.9
51513.8647	0.1821	-124.2	-2.5	190.0	-4.0
51513.8779	0.2016	-126.9	2.5	200.3	-1.2
51513.8887	0.2176	-130.5	3.3	204.4	-1.4
51513.9011	0.2359	-133.6	3.2	205.2	-3.5
51513.9118	0.2517	-133.4	4.1	208.4	-0.9
51513.9238	0.2694	-128.8	7.4	206.8	-1.3
51513.9346	0.2854	-123.6	9.6	207.2	2.1
51524.7788	0.3022	-116.4	11.7	202.9	2.7
51524.7912	0.3205	-116.4	4.1	198.1	5.3
51524.8062	0.3427	-103.1	5.4	187.5	6.3
51524.8169	0.3585	-99.0	-0.9	170.1	-1.1
51524.8289	0.3762	-82.3	2.7	174.9	16.5
51524.8397	0.3922	-73.0	-1.2	163.9	18.3
OU Ser					
50852.8082	0.2676	-108.4	-4.0	163.1	-5.6
50852.8133	0.2848	-109.9	-6.2	162.1	-2.5
50852.8184	0.3020	-103.6	-1.1	159.1	1.3
50852.8251	0.3246	-106.1	-5.8	145.1	0.2
50852.8302	0.3417	-98.1	0.0	130.8	-1.5
50852.8354	0.3593	-97.0	-1.5	118.3	1.2
50852.8417	0.3805	-89.4	2.4	101.1	5.4
50852.8472	0.3990	-91.3	-3.2	82.4	7.6
50852.8525	0.4169	-91.2	-6.9	...	...
50852.8586	0.4374	...	...	...	...
50852.8638	0.4550	...	...	...	...
50852.8690	0.4725	...	...	...	...
50852.8757	0.4951	...	...	...	...
50852.8809	0.5126	...	...	...	...
50852.8860	0.5298	...	...	...	...
50852.8926	0.5520	...	...	...	...
50852.8978	0.5695	...	...	...	...
50852.9030	0.5871	...	...	...	...
50852.9090	0.6073	...	...	...	...
50852.9143	0.6251	-47.6	-12.2	-243.8	-13.9
50852.9196	0.6430	-37.0	-4.7	-248.1	-0.8
50852.9273	0.6689	-34.2	-5.6	-275.6	-7.0
50852.9324	0.6861	-28.8	-2.1	-279.3	0.4
50852.9375	0.7033	-29.1	-3.9	-291.7	-3.4
50852.9436	0.7239	-25.7	-1.7	-293.4	1.8
50852.9486	0.7407	-25.7	-2.1	-292.8	5.1
50852.9537	0.7579	-26.0	-2.5	-294.7	3.3
50852.9597	0.7781	-24.5	-0.4	-293.9	0.8
50852.9648	0.7953	-26.1	-1.0	-287.5	1.4
50852.9699	0.8125	-27.1	-0.5	-280.8	-0.3
50852.9765	0.8347	-26.2	2.9	-263.5	2.4
50852.9817	0.8523	-26.7	4.9	-252.2	-0.6
50852.9868	0.8694	-31.6	2.8	-233.4	2.0
50858.8758	0.7135	-26.4	-1.9	-291.0	1.2

TABLE 1. (continued)

HJD-2,400,000	Phase	V_1	ΔV_1	V_2	ΔV_2
50858.8817	0.7334	-28.6	-4.9	-299.8	-2.8
50858.8878	0.7539	-28.6	-5.1	-297.1	1.1
50858.8945	0.7765	-21.4	2.6	-291.2	3.9
50858.9011	0.7987	-27.7	-2.3	-291.1	-3.7
50858.9077	0.8210	-32.6	-5.1	-276.0	-0.6
50858.9151	0.8459	-30.8	-0.2	-265.3	-8.2
50858.9216	0.8678	-27.0	7.1	-236.7	0.3
50858.9279	0.8890	-36.4	1.6	-210.4	4.1
50896.9083	0.8709	-32.8	1.8	-230.0	3.9
50896.9155	0.8951	-26.3	12.9	-202.3	5.2
50896.9231	0.9208	...	...	...	...
50896.9320	0.9507	...	...	...	...
50896.9392	0.9750	...	...	...	...
50930.8318	0.1823	-102.4	-1.3	148.1	-1.2
50930.8380	0.2032	-97.7	5.2	160.2	0.1
50930.8442	0.2240	-103.7	0.4	164.4	-2.6
50930.8520	0.2503	-106.1	-1.4	173.0	2.8
50930.8583	0.2716	-105.0	-0.7	161.0	-7.0
50930.8647	0.2931	-99.6	3.6	165.5	3.9
50930.8720	0.3177	-102.9	-1.8	152.3	3.0
50930.8778	0.3373	-92.5	6.2	140.9	5.1
50930.8840	0.3582	-93.8	1.9	123.1	5.0
50930.8909	0.3814	-94.2	-2.6	104.0	9.3
50930.8968	0.4013	-91.2	-3.5	...	...
50930.9024	0.4202	-84.7	-1.1	...	...
50940.8035	0.7837	-20.9	3.5	-289.3	3.8
50940.8098	0.8049	-29.8	-3.9	-280.1	4.4
50940.8184	0.8339	-24.2	4.8	-259.7	6.8
50949.7573	0.9552	...	...	...	...
50949.7632	0.9750	...	...	...	...
50949.7691	0.9949	...	...	...	...
50949.7750	0.0148	...	...	...	...
50949.7823	0.0394	...	...	...	...
50949.7884	0.0600	...	...	...	...
50949.7942	0.0795	...	...	...	...
50949.7999	0.0987	...	...	...	...
50949.8065	0.1210	-90.5	1.5	98.3	1.0
50949.8124	0.1408	-95.5	-0.0	122.1	4.9
50949.8182	0.1604	-100.2	-1.8	137.7	3.7
50949.8250	0.1833	-101.1	0.1	150.2	0.3
50949.8308	0.2028	-103.9	-1.0	159.7	-0.2
50949.8367	0.2227	-106.4	-2.3	161.2	-5.5

^aThe data which have been given half weight in the orbital solutions.

^bThe data for UX Eri kindly provided by Dr. H. Duerbeck.

Notes to Table 1.

Velocities are expressed in km s^{-1} . Observations leading to entirely unseparable broadening- and correlation-function peaks are marked by the “no-data” symbol (...); these observations may be eventually used in more extensive modeling of broadening functions.

TABLE 2. Spectroscopic orbital elements of the third ten of close binary systems

Name Sp. type	Type	Other names	γ	K_1 K_2	ϵ_1 ϵ_2	$T_0 - 2,400,000$ (O–C) and [E]	P (days) assumed	q = m_2/m_1
CN And F5V	A/EB	BD+39°59	–24.86(1.45)	87.53(1.11) 224.35(2.84)	6.10 18.03	50850.2790(22) –0.0137 [327]	0.462793	0.390(33)
HV Aqr F5V	A	BD–3°5183	–3.77(1.25)	42.41(1.44) 293.35(2.11)	4.83 11.46	50859.3453(8) ... ^a	0.374460	0.145(50)
AO Cam G0V	W	BD+52°826	–13.17(0.83)	250.50(1.05) 103.51(1.28)	6.47 5.50	50960.1131(5) –0.0121 [13732]	0.329905	2.420(11) 1/q=0.413
YY CrB F8V	A	HD 141990 HIP 77598	–4.58(1.02)	68.07(1.54) 279.85(1.55)	7.79 14.13	50955.8711(6) –0.0053 [6521]	0.376565	0.243(23)
FU Dra F8V	W	HIP 76272	–11.38(1.14)	280.76(2.09) 70.38(1.84)	8.68 7.51	50866.2777(8) –0.0086 [7714]	0.306718	3.989(30) 1/q=0.251
RZ Dra A6V	A/EB	HIP 90092	10.68(1.76)	96.85(1.72) 244.47(3.54)	7.53 22.22	51035.4103(28) –0.0165 [39186]	0.550876	0.396(37)
UX Eri F9V	A	BD–7°553 HIP 14699	12.79(1.09)	91.75(1.55) 245.76(1.86)	6.81 10.09	50416.5587(14) +0.0030 [–78]	0.445279	0.371(21)
RT LMi F7V:	A		–10.30(1.74)	95.34(1.67) 260.17(3.59)	7.28 15.68	51261.1008(16) –0.0016 [41056]	0.374918	0.366(38)
V753 Mon A8V	W	HD 54975 HIP 34684	38.58(0.83)	176.07(0.97) 170.76(0.97)	8.84 6.80	51188.0809(15) –0.0249 [3970]	0.677049	1.031(9) 1/q=0.970
OU Ser F9/G0V	A	HD 136924 HIP 75269	–64.08(0.41)	40.59(0.59) 234.24(0.69)	3.80 5.97	50901.9916(3) +0.0026 [8093]	0.296764	0.173(17)

^aThe two available ephemerides for moments of primary minima of HV Aqr give discordant results and do not provide a clear guidance to which eclipse our T_0 applies. It has been assumed that the system is A-type on the basis of the light curve data; see the text.

Notes to Table 2.

The convention of naming the components is that the subscript 1 designates the component which is eclipsed at the deeper minimum and is therefore the hotter one. The standard errors of the circular solutions in the table are expressed in units of last decimal places quoted; they are given in parantheses after each value. For example, the last table entry for q , 0.173(17) should be interpreted as 0.173 ± 0.017 . The average radial velocities (γ), the velocity amplitudes (K_i) and the standard unit-weight errors of the solutions (ϵ) are expressed in km s^{-1} . The calculated moments of primary minima are given by T_0 while their (O–C) deviations (in days) have been calculated from the most recent available ephemerides, as given in the text, using the assumed periods and the number of epochs given by [E].